




Reagents for Electrophilic Aromatic Substitution:

(Klein Sections 18.1-18.6, generate E^+)

Preparation of Diazonium Salt (ArN_2^+):

Reagents for Sandmeyer Reactions:


(Klein Section 22.11, react with ArN_2^+)


Apply ArN_2^+ in synthesis (Try text problem 22.26) 18-12



* can't be done
with EAS alone!


Ex. Synthesize following TM from benzene or toluene.



Example: Transform

*Note: can't be made
via Friedel-Crafts
Alkylation (NO_2 is EWG, so N/R)



18-13





Reactions of Aromatic substituents (17.6) 18-14

↳ groups attached / side chains

a) Reduction Rxns (review)




b) Oxidations of Arenes




* benzylic [ox] even breaks C-C bonds!



* benzylic carbon can't be quaternary



Recall: Other benzoic acid preps (20.4)



c) benzylic halogenation (10.5)


18-15



* easy to introduce a benzylic LG!

→ radical stabilities: benzylic/allylic/ $3^\circ > 2^\circ > 1^\circ > \text{methyl}$

→ NBS is also a source of Br.



d) benzylic substitutions


* S_N1 and S_N2 are both great with benzylic LG's



→ S_N2 favored over $E2$


→ benzylic TS. is stable

Ex. Transform






Skill Builder 17.4

Nucleophilic Aromatic Substitution, S_NAr (18.13, 18.15) 18-16




- * o/p EWGs are needed to stabilize $\ominus$ intermediate
- * ring is $E^\oplus$ so ring is _____ toward S_NAr by EWG
- * LG ability: $F > Cl > Br > I$



* SkillBuilder 18.7 *

Note: skip benzyne (18.14) FYI see Educator 52:46-59:10



California State Polytechnic University, Pomona

Organic Chemistry II, CHM 3150, Dr. Laurie S. Starkey

Ch. 17-18 Summary (Klein): Aromaticity & Aromatic Substitution Reactions

I. Benzene (17.3)

- A) resonance stabilized (confirmed ΔH hydrogenation), 3-D sketch

II. Aromaticity

- A) definition and rules (17.1, 17.4) **SkillBuilder 17.2**

- i) cyclic and planar; contiguous p orbitals
- ii) $(4n + 2)$ electrons in p orbitals (Hückel's rule, $n = 0, 1, 2$, etc.)
- iii) Molecular Orbital (MO) theory to explain aromatic stability

- B) common aromatic compounds (furan, naphthalene, etc.) (17.5) **SkillBuilder 17.3**

- C) nomenclature: *ortho* (*o*-), *meta* (*m*-), *para* (*p*-) positions, common names (17.2) **SB 17.1**

- D) special topics: buckminsterfullerene (C_{60} , Bucky Ball) as a "new" form of carbon

III. Electrophilic Aromatic Substitution (EAS) (18.2 – 18.6)

- A) mechanism: formation of E^+ (one or more steps) addition of E^+ (slow step), loss of H^+

IV. EAS on substituted benzenes (18.7 – 18.11) **SkillBuilder 18.1, SkillBuilder 18.3**

- A) three categories of substituents (18.10)

- i) electron-donating groups (EDG, 18.7)
- ii) electron-withdrawing groups (EWG, 18.8)
- iii) halogens ($X = Cl, Br$) (18.9)

- B) reactivity

- i) EDG are activating (electron-rich ring, good Nu:)
- ii) EWG/X are deactivating (electron-deficient ring, poor Nu:)

- C) regioselectivity (for *ortho*, *para* directors, *para* is usually major due to sterics)

- i) regioselectivity can be explained by looking at resonance forms for:
 - a) starting material (Nu:) electron density (in certain cases only)
 - b) carbocation intermediate stability (can be used for all cases)
- ii) EDG/X are *ortho*, *para* directors (because they stabilize adjacent carbocations)
- iii) EWG are *meta* directors (because EWG's destabilize adjacent carbocations)

- D) directing power for disubstituted benzenes (18.11) **SkillBuilder 18.2**

V. Electrophiles for EAS

- A) $-X$ ($Br_2/FeBr_3$ or $Cl_2/FeCl_3$) (18.2)

- B) $-NO_2$ (HNO_3/H_2SO_4) (18.4)

- C) $-SO_3H$ (SO_3/H_2SO_4), reaction is reversible (18.3) **SkillBuilder 18.4**

- D) $-R$ (Friedel-Crafts Alkylation, via carbocation which can rearrange) (18.5)

- i) $RX + AlCl_3$ or $ROH + HA$ or $ROH + BF_3$ or alkene + HA

- E) $-COR$ (Friedel-Crafts Acylation, $RCOX/AlCl_3$) (18.6)

- i) can then reduce carbonyl to give desired 1° alkyl side chain

VI. Diazonium Salts (ArN_2^+ , 22.10, 22.11)

- A) two-step preparation from $ArNO_2$

- i) reduce nitro to amine: $-NO_2 \rightarrow -NH_2$ (H_2/Pd or $Fe/Zn/Sn$ and HA)

- ii) add nitrous acid $-NH_2 \rightarrow -N_2^+$ ($HONO = NaNO_2/HCl$)

- B) uses: replace its good LG (N_2) with the following groups

- i) halogens: $-Cl$ ($CuCl$), $-Br$ ($CuBr$), $-I$ (KI), $-F$ (HBf_4)

- ii) cyanide: $-CN$ ($CuCN$)

- iii) hydrogen: $-H$ (H_3PO_2)

- iv) hydroxyl $-OH$ (H_2O/H_2SO_4)

VII. Aromatic Synthesis (18.12) **SkillBuilders 18.4, 18.5, 18.6**

VIII. Nucleophilic Aromatic Substitution (S_NAr , 18.13, 18.15) **SkillBuilder 18.7**

- A) two-step mechanism: add Nu:, eject LG (Addition-Elimination)

- B) EWG substituents ($-NO_2$) required

IX. Reactions of Benzylic Carbons (17.6) **SkillBuilder 17.4**

- A) oxidation (Na_2CrO_7 or $KMnO_4$) to oxidize benzylic carbons $\rightarrow$ benzoic acids

- B) substitution (S_N1 , S_N2) and free-radical halogenation ($Br_2/h\nu$)

SKIP: Spectroscopy (17.8), Benzyne "Elimination-Addition" (18.14), [ox] of $ArOH$ (12.12)